

**EFFECTS OF DIRAC'S NEGATIVE ENERGY
SEA ON QUANTUM NUMBERS**

R. Jackiw

*Center for Theoretical Physics
Laboratory for Nuclear Science
and Department of Physics
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139 U.S.A.*

Dirac Prize Lecture
Trieste, Italy – March 1999

THE DIRAC PRIZE COMMITTEE cited my work on fractional charges and on chiral anomalies; therefore, I shall discuss these two topics. As with all true and deep physical effects, there are many ways of arriving at the results. It is particularly appropriate here today that one route towards understanding both fractional charges and chiral anomalies delves into Dirac's negative energy sea. This is especially provocative, because usually we think of Dirac's negative energy sea as an unphysical construct, invented to render quantum field theory physically acceptable by hiding – that is, by renormalizing – negative energy solutions. But I am suggesting that in fact physical consequences can be drawn from Dirac's construction.

First let me set the stage for the discussion.

Quantum physics has taught us that a physical observable need not be a quantity with arbitrary magnitude. Because it is the eigenvalue of a linear Hermitian operator, it will in general be quantized. This is not the case in classical physics where most observables, like angular momentum and energy, are continuously varying and can take on any value. On the other hand, even in classical physics, there are concepts that are intrinsically integral – for example particle number of conserved particles – and one expects that the integrality will be preserved in the quantum theory, that is, eigenvalues of the relevant operator – the number operator in our example – are expected to be integers. Quantization of eigenvalues is most easily attained when the operator is a generator of a compact, non-Abelian group, like angular momentum generating $SO(3)$ rotations. However, the number operator frequently generates only Abelian transformations with no group-theoretic quantization.

Closer examination of the number operator in a theory with second quantized fermions raises doubts that it will in fact possess only integer eigenvalues. The problem derives from Dirac's negative energy sea, which must be filled to define the vacuum. This involves an infinite number of "particles". Since the number of any further particles must be measured relative to this infinity, there may very well emerge a non-integral answer. Nevertheless, it had been believed that various renormalization procedures, like normal ordering, can unambiguously ensure integrality of the eigenvalue. Therefore, it was a surprise when Rebbi and I established about twenty years ago that fermions moving in the field of a topologically non-trivial soliton (kink in one spatial dimension, vortex in two, monopole in three) possess non-integer eigenvalues for their number operator.¹ It is perhaps even more surprising that this peculiar effect has a physical realization in properties of actual condensed matter systems – polyacetylene being the standard example.² Here, I shall describe this to you, first in a general, formal way, and then in a physically intuitive language appropriate to polyacetylene.

We wish to second quantize fermions moving in a static background that generically is described by φ . Fermion dynamics is governed by a Dirac Hamiltonian $H(\varphi)$. Two different backgrounds are envisioned: one is appropriate for the vac-

uum sector of the theory φ_v , the other for the soliton φ_s . For example, φ may be a condensate field that takes a homogenous value in the vacuum sector and a topologically nontrivial profile in the soliton sector.

Second quantization proceeds by computing the energy eigenvalues and eigenfunctions of $H(\varphi)$, which possesses both positive and negative energy eigenstates, and “filling” the negative energy sea. The number density of the soliton ground state is given by

$$\rho(\mathbf{x}) = \int_{-\infty}^{0^-} dE (|\Psi_E(\mathbf{x})|^2 - |\psi_E(\mathbf{x})|^2) \quad (1)$$

where the integration, which also includes summation over discrete levels, extends over all negative energy states, since they are filled in the vacuum. Here Ψ_E is the energy eigenfunction in the presence of the soliton and ψ_E is the eigenfunction in the vacuum sector:

$$H(\varphi_s)\Psi_E = E\Psi_E, \quad H(\varphi_v)\psi_E = E\psi_E \quad . \quad (2)$$

In (1) the contribution from the vacuum sector is subtracted; the soliton charge density is renormalized, so that it is measured relative to the vacuum. The fermion number of the soliton ground state is the spatial integral of ρ

$$N_F = \int d\mathbf{x} \rho(\mathbf{x}) \quad . \quad (3)$$

A very beautiful aspect of the theory is that one can evaluate (1) and (3) by general methods, which bypass solving the eigenvalue problem (2) explicitly. Rather, one uses spectral sum rules whose form is dictated by general features of the Hamiltonian, in particular by the topological properties of the background φ and of the space $\{\mathbf{x}\}$. While these methods are powerful, they are also technical, requiring much mathematical knowledge, so I shall not present them here. However, when the Hamiltonian possesses one further property, which I shall now describe, the sum rules become trivial, and the result for N_F is immediate.

We assume further that $H(\varphi)$ possesses a conjugation symmetry, taking positive energy states into negative energy states and *vice versa*; that is, we assume there exists an operator \mathcal{C} that anticommutes with $H(\varphi)$: $\mathcal{C}^{-1}H\mathcal{C} = -H$. One consequence of this is that the number density at E is an even function of E : $|\Psi_E|^2 = |\Psi_{-E}|^2$, $|\psi_E|^2 = |\psi_{-E}|^2$. A less obvious consequence is that in the soliton sector, there are always normalizable, discrete zero-energy modes

$$H(\varphi_s)u_0 = 0, \quad \int d\mathbf{x} |u_0(\mathbf{x})|^2 = 1 \quad . \quad (4)$$

This fact may be seen by explicit solution of the eigenvalue problem, but it also follows from a general mathematical argument, called *index theory*, which ensures that the Dirac operator has normalizable zero-energy modes in the presence of a topologically nontrivial background.

We are now in a position to evaluate (1) and (3). First, we use completeness of the eigenfunctions in the soliton and vacuum sectors

$$\int_{-\infty}^{0^-} dE |\Psi_E(\mathbf{x})|^2 + \int_{0^+}^{\infty} dE |\Psi_E(\mathbf{x})|^2 + |u_0(\mathbf{x})|^2 - \int_{-\infty}^{\infty} dE |\psi_E(\mathbf{x})|^2 = 0 \quad . \quad (5)$$

The zero-energy mode in the soliton sector has been explicitly separated; we assume there is just one. (In the vacuum sector there are none.) Then, use of the conjugation symmetry allows equating the positive energy integrals with the negative energy ones, and converts (5) into an evaluation of (1)

$$\int_{-\infty}^{0^-} dE (|\Psi_E(\mathbf{x})|^2 - |\psi_E(\mathbf{x})|^2) = -\frac{1}{2}|u_0(\mathbf{x})|^2 \quad . \quad (6)$$

The spatial integration that determines N_F is trivial since the zero mode is normalized:

$$N_F = -\frac{1}{2} \quad . \quad (7)$$

The conclusion is that the soliton vacuum, defined with the zero mode empty, carries fermion number $-\frac{1}{2}$; of course, when the zero mode is filled, this fermion number is $+\frac{1}{2}$! The fermion number assignment of $\pm\frac{1}{2}$ for two states degenerate in energy is the only possible one consistent with a conjugation-odd fermion number operator.

Two comments should be made in connection with this very elementary derivation of our surprising result.

- (i) The above evaluation concerns the *expected* value of the second-quantized, field theoretic number operator, \hat{N}_F . However, one can show, by expanding the second quantized field in terms of creation and annihilation operators in the presence of the soliton, that in fact the *eigenvalues* are $\pm\frac{1}{2}$.
- (ii) We have viewed the soliton as an external field. In a complete description, one must take the soliton's quantum dynamics into account. Necessarily there will occur spontaneous symmetry breaking in the vacuum sector. Calculations in the full theory can be carried out by Monte-Carlo methods, or by analytic techniques of the Born-Oppenheimer variety.

The three ingredients necessary for fermion number fractionization – spontaneous symmetry breaking, solitons and fermions – come together in a description of a physical system, polyacetylene. This is a one-dimensional array of carbon atoms that can form one of two degenerate ground states. The degeneracy arises from a spontaneous breaking of the right-left symmetry (Peierls instability) and manifests itself in an alteration of the bonding pattern, as illustrated in Fig. 1.

A microscopic Hamiltonian for the system has been proposed by Su, Schrieffer and Heeger (SSH).² In the continuum and infinite volume limit, electron transport is governed by a Dirac-type Hamiltonian in one spatial dimension:

$$H(\varphi) = \sigma_3 \hat{p} + \sigma_1 \varphi(x), \quad \hat{p} = \frac{1}{i} \frac{d}{dx} \quad (8)$$

$$\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad .$$




Fig. 1: Polyacetylene consists of a linear chain of carbon atoms (dots). The equally spaced configuration (O) possesses a left-right symmetry, which however is energetically unstable. Rather in the ground states the carbon atoms shift a distance μ to the left or right, breaking the symmetry and producing two degenerate ground states (A,B). (The drawing is not to scale; the shift is only a few percent of the total bond length.) A soliton (S) is a defect in the alteration patterns; it provides a domain wall between configurations (A) and (B).

Here, $\varphi(x)$ is the phonon field; it measures the displacement of the carbon atom from its equilibrium position. The matrix structure of the above Hamiltonian does not arise from spin. In the SSH description, electron-electron interactions are ignored and spin is a passive label; the Hamiltonian in (8) describes separately spin up and spin down electrons. Rather, the Dirac-like matrix form for H arises through a linearized approximation and the two-component wavefunctions that are eigenmodes of H refer to the right-moving and left-moving electrons with momentum $\pm|p|$. The filled negative energy states are the valence electrons, while the conduction electrons populate the positive energy states.

In the SSH model, $\varphi(x)$ is determined self-consistently by the phonon's dynamics, and in the lowest (vacuum) states $\varphi(x)$ takes the uniform values $\pm\mu$, as illustrated in Fig. 1. The corresponding spectrum of (8) exhibits a gap.

In addition to the two ground states, where the phonon field takes a constant value, there exist stable excited states where $\varphi(x)$ assumes a kink shape, which interpolates as x passes from $-\infty$ to $+\infty$, between the vacuum configurations $-\mu$ and $+\mu$. This is the soliton, and it describes a defect in the alteration pattern, as is also exhibited in Fig. 1.

The Hamiltonian in (8) admits a conjugation symmetry: $\mathcal{C} = \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ anticommutes with H – this is ordinary charge conjugation invariance in the absence of Coulomb interactions. The zero eigenvalue problem is easily solved with a kink background: there is one normalizable zero-energy solution. Thus, our general analysis predicts that fermion number, here coinciding with charge, fractionizes to $\pm\frac{1}{2}$ in the one-soliton state.

This result may also be seen pictorially. When two solitons are inserted into the ground state (B), the bonding pattern is depicted in Fig. 2. Note that the number of bonds in the two-soliton state is one fewer than in the ground state. If the two solitons are now separated far apart, so that they act independently, the quantum numbers of the missing bond must be shared between the two states, and that is how the fraction $\frac{1}{2}$ arises.




Fig. 2: With two solitons (SS) inserted in vacuum (B), the number of bonds \approx electrons between the sites of the defects decreases from five to four.

Of course, for actual physical samples, where both the volume and the separation between defects are finite, the non-integer fermion number is only an expectation value for the operator \hat{N}_F , and there are corrections that vanish in the infinite limit. The important point is that the variance $\langle \hat{N}_F^2 \rangle - \langle \hat{N}_F \rangle^2$ also vanishes in the limit. This is to be contrasted to the uninteresting situation of, say, an electron circulating about two nuclei. When the nuclei are far apart, the expected value for electron number near each nucleus is $\frac{1}{2}$, plus small corrections. However, the variance remains $\frac{1}{4}$ for infinite separation, which shows that this fraction never becomes an eigenvalue.

The concept of fractional quantum numbers has now extended beyond soliton systems – for example the theory of the quantum Hall effect makes use of the idea, even though dynamical details are quite different from the above example.

While fractional quantum numbers were first seen in relativistic field theory, thus far they have not played any experimentally verified role in particle physics. Nevertheless, it is curious that an effect which in principle is physical, and has been observed in condensed matter systems, should arise from distortions in the negative energy sea, which for particle physicists is an unphysical construct, in contrast to the condensed matter application, where negative energy states correspond to physical quantities – the valence electrons.

In particle physics there is another, physically realized circumstance where the Dirac negative energy sea modifies symmetry behavior of fermions. This is the chiral anomaly phenomenon whereby the axial vector current $i\bar{\psi}\gamma^\mu\gamma^5\psi$, which is conserved for free massless Dirac fermions, ceases to be conserved when the massless fermions are quantized and made to interact with a gauge field, even though the interaction appears to be chirally invariant.³ Indeed, for unquantized Dirac fields

the chiral invariance ensures conservation of the unquantized axial vector current. This disappearance after quantization of chiral symmetry is usually associated with infinities that plague relativistic quantum field theory: the infinities must be regulated and renormalized, but there is no chirally invariant regulator procedure. However, a more directly physical discussion of the anomaly phenomenon may be given, which shows that in fact it is the filling of the negative energy sea that breaks the chiral symmetry.

Let me first state the essential puzzle of the chiral anomaly. In spite of profound differences between the classical and quantal description of physical phenomena, it was generally believed that symmetry properties and conserved constants of motion transcend the classical/quantal dichotomy: when a classical model possesses symmetries and supports constants of motion, one expects that quantization preserves the symmetries, so that conserved quantities – now promoted to quantum operators – remain time-independent, that is, they commute with the Hamiltonian operator. But as observed thirty years ago by Bell and me,³ and also independently by Adler,³ this need not be so.

A simple instance of quantum mechanical violation of symmetry is encountered by considering a massless Dirac fermion moving in a background gauge field A_μ . The dynamics is governed by a Lagrangian, which splits into separate right and left parts – this separation is a manifestation of chiral symmetry:

$$\begin{aligned} \mathcal{L} &= \bar{\psi}(i \not{\partial} - e \not{A})\psi \\ &= \bar{\psi}_+(i \not{\partial} - e \not{A})\psi_+ + \bar{\psi}_-(i \not{\partial} - e \not{A})\psi_- \\ \psi &= \psi_+ + \psi_-, \quad \psi_\pm = \frac{1}{2}(1 \pm i\gamma_5)\psi \quad . \end{aligned} \tag{10}$$

In the first-quantized theory, where ψ is a wavefunction and $\bar{\psi}\gamma^\mu\psi$ is a probability current, the separate right and left currents are conserved, and the separate probabilities $\int d\mathbf{x} \psi_\pm^\dagger \psi_\pm$ are time-independent. In the second quantized theory, where ψ becomes an operator, the anomaly phenomenon renders the separate right and left currents no longer conserved, and the right and left charges are not time-independent. Nevertheless, the sum of right and left currents – the vector current – is conserved, while the divergence of the difference between the right and left currents – the axial vector current – is nonzero owing to the anomaly. Our task then is to understand what causes the separate nonconservation of left and right currents even though there is no coupling between the two apparent in (10).

Evidently, the problem derives from the second quantization procedure, hence we implement it. We set A^0 to zero, find the eigenmodes of the Hamiltonian in the background field \mathbf{A} , and define the second quantized vacuum by filling the negative energy modes, leaving the positive energy modes empty. The background \mathbf{A} is chosen in a specific functional form so that the anomaly is nonvanishing. This requires \mathbf{A} be time-dependent, but we chose a potential constant in time and space and model the time variation by an adiabatic change $\mathbf{A} \rightarrow \mathbf{A} + \delta\mathbf{A}$.

The simplest model to study is two-dimensional and Abelian – two-dimensional massless quantum electrodynamics.⁴ The Dirac matrices are 2×2 and ψ is a two-

component spinor.

$$\begin{aligned} \gamma^0 &= \sigma^1, & \gamma^1 &= i\sigma^2, & \gamma_5 &= i\gamma^0\gamma^1 = -i\sigma^3 \\ \psi_+ &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \psi, & \psi_- &= \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \psi. \end{aligned} \quad (11)$$

The axial current possess an anomalous divergence proportional to $\epsilon^{\mu\nu} F_{\mu\nu} \propto \partial_t A$ (since $A_0 = 0$ and \mathbf{A} has the single spatial component $A^1 \equiv A$),⁵ which we wish to understand. The eigenmodes to be second quantized satisfy a one-dimensional Dirac equation,

$$H\psi_E = \alpha(\hat{p} - eA)\psi_E = E\psi_E, \quad \alpha = -\sigma^3 \quad (12)$$

where A is constant. They are given by

$$\begin{aligned} \psi_+ &= \begin{pmatrix} e^{ipx} \\ 0 \end{pmatrix} \text{ with } E = -p + eA \\ \psi_- &= \begin{pmatrix} 0 \\ e^{ipx} \end{pmatrix} \text{ with } E = p - eA \end{aligned} \quad (13)$$

Second quantization is performed by filling the negative energy sea. For $A = 0$, the energy-momentum dispersion is depicted in Fig. 3, where the right-hand branch corresponds to fermions of one chirality, and the left-hand branch to those of the other chirality. The negative energy states are filled, as indicated by the filled circles; the positive energy states are empty, as indicated by the empty circles. As A increases from 0 to δA , empty states in the right-hand branch acquire negative energy, while filled states of the left-hand branch become positive energy states; that is, there is a net production of right-handed antiparticles and left-handed particles; see Fig. 4. So the separate right and left charges are not conserved, but their sum is. Put in another way, the separation between positive and negative energy states of definite chirality cannot be achieved gauge-invariantly, since changing A from 0 to a constant δA is a gauge transformation, yet particles are produced.




Fig. 3: Energy-momentum dispersion at $A = 0$. Empty circles are empty states; filled circles are filled states.




Fig. 4: Energy-momentum dispersion at $A = \delta A > 0$. The energy shift produces negative energy empty states and positive energy filled states.

We thus see that the negative energy sea is responsible for nonconservation of chirality even though the dynamics is chirally invariant. This effect was called *anomalous* because its discovery was a surprise.³ However, a better name might be *quantum mechanical symmetry breaking* – a symmetry-breaking mechanism, which like Heisenberg's *spontaneous symmetry breaking* attributes physical asymmetry to the vacuum state and not to the dynamics. Here, however, unlike in Heisenberg's case, it is not vacuum degeneracy but the very definition of the vacuum that is responsible. Once again we must assign physical reality to Dirac's negative energy sea, because it produces the chiral anomaly, whose effects are experimentally observed, principally in the decay of the neutral pion to two photons, but there are other physical consequences as well.⁶

The two phenomena that I have described show that Nature seems to know and make use of what at first appears to be a defect of a quantum field theory, and which is usually “renormalized away”. Remarkably, the infinities in the formalism give rise to finite and physical effects. One may quite appropriately call this an example of “The Unreasonable Effectiveness of Quantum Field Theory in Physics”.

In my presentation of fractional charge and of the chiral anomaly I have used concepts that are primitively physical, employing little analysis beyond drawing pictures and counting. But to the same end a most sophisticated mathematical discussion can also be given, wherein zero modes are controlled by various index theorems (Atiyah-Singer, Callias), fractional charges are related to Atiyah, Patodi, Singer spectral flows, and chiral anomalies are identified with Chern-Pontryagin densities. This confluence between physics and mathematics, which was brought about by physicists' research on the topics that I described, seeded an interaction between these two disciplines, which still flourishes and today fuels the current string/M-theory program.

But in sharp contrast with the above, contemporary mathematical discussions within physics do not as yet have experimentally observed correlatives. These days research follows very closely Dirac's dictum:

The most powerful method of advance [in physics] . . . is to employ all the resources of pure mathematics in attempts to perfect and generalize the mathematical formalism that forms the existing basis of theoretical physics, and . . . to try to interpret the new mathematical features in terms of physical entities.
— *P.A.M. Dirac*

Early in my career I heard him say this at a seminar devoted to his research history, and I was very inspired. However, my subsequent work with physics and mathematics makes me feel that Dirac's suggestion is too radical, while a quote by Yang more accurately describes my own experience of mathematical physics:

Physics is not mathematics, just as mathematics is not physics. Somehow nature chooses only a subset of the very beautiful and complex and intricate mathematics that mathematicians develop, and that precise subset is what the theoretical physicist is trying to look for.
— *C.N. Yang*

REFERENCES

1. R. Jackiw and C. Rebbi, *Phys. Rev. D* **13**, 3398 (1976); for a review, see A. Niemi and G. Semenoff, *Phys. Reports* **135**, 99 (1986).
2. W.-P. Su, J.R. Schrieffer and A. Heeger, *Phys. Rev. Lett.* **42**, 1698 (1979) and *Phys. Rev. B* **22**, 2099 (1980).
3. J.S. Bell and R. Jackiw, *Nuovo Cim.* **60**, 47 (1969); S. Adler, *Phys. Rev.* **177**, 2426 (1969).
4. J. Schwinger, *Phys. Rev.* **128**, 2425 (1962).
5. K. Johnson, *Phys. Lett.* **5**, 253 (1963).
6. For extensive discussion of the anomaly phenomena see S. Treiman, R. Jackiw, B. Zumino and E. Witten, *Current Algebra and Anomalies*, (Princeton University Press/World Scientific, Princeton, NJ/Singapore, 1985) or S. Adler in *Lectures on Elementary Particles and Quantum Field Theory*, S. Deser, M. Grisaru, and H. Pendleton, eds. (MIT Press, Cambridge, MA, 1970).